arXiv:1609.03690v1 [cs.IT] 13 Sep 2016

Construction of Gray maps for groups of order 16

Ko Sakai[1] and Yutaka Sato[2]
[1]Faculty of Pure and Applied Sciences,
[2]Graduate School of Pure and Applied Sciences,
University of Tsukuba, Tsukuba city, Japan

September 14, 2016

Abstract. In this paper we consider Type 1 Gray maps and Type 2 Gray maps for groups of order 16. First, we confirm that we can construct Type 1 Gray maps for all groups of order 16, and we actually construct them. Next, we confirm that we can construct Type 2 Gray maps for several groups of order 16, and we construct all such maps. Finally we reveal why the other groups do not admit Gray maps.

1 Introduction

Reza Sobhani constructed two classes of Gray maps, called Type 1 Gray map and Type 2 Gray map, for finite p-groups G in [1]. Type 1 Gray maps are constructed based on a Gray map for a maximal subgroup H of G. In contrast, Type 2 Gray maps are constructed, if G is a semidirect product of two finite p-group H and K, both H and K admit Gray maps and the corresponding homomorphism $\psi : H \to \mathrm{Aut}(K)$ is compatible with the Gray map for K in the sense that for all $h \in H$ we have $w_H(\theta_2(k)) = w_H(\theta_2(\psi_h(k)))$. Type 2 Gray maps have feature which they generally generate shorter codes than Type 1 Gray maps. However, we can construct for only 6 groups among all the groups of order 16.

Marcel Wild gave complete classification of the groups of order 16 based on several elementary facts. He examined them as extensions of groups of order 8 by the cyclic group of order 2.

In this paper we consider Type 1 Gray maps and Type 2 Gray maps for groups of order 16 in the same line as Sobhani. We assume that information is encoded in an alphabet Q with q distinct symbols.

2 Preliminaries

In this section we assume that G is a finite 2-group of order 2^m.

2.1 Hamming-distance, Hamming-weight and Gray map

Definition 1 For any two elements $\mathbf{u} = (u_1, u_2, \ldots, u_n)$ and $\mathbf{v} = (v_1, v_2, \ldots, v_n)$ in Q^n, the *Hamming-distance* between $\mathbf{u}$ and $\mathbf{v}$ is defined by,

$d_H(\mathbf{u}, \mathbf{v}) \stackrel{\text{def.}}{=} |\{i \mid 1 \le i \le n, u_i \ne v_i\}|$.

Definition 2 The *Hamming-weight* of $\mathbf{u}$ is defined by,

$w_H(\mathbf{u}) \stackrel{\text{def.}}{=} |\{i \mid 1 \le i \le n, u_i \ne 0\}|$.

We review the definition and a lemma on a Gray map in [1].

Definition 3 A map $\phi : G \to \mathbb{Z}_2^n$ is said to be a *Gray map*, if the following properties hold,
(1) The map $d_\phi : G \times G \to \mathbb{N} \cup \{0\}$ defined by $d_\phi(a, b) = w_H(\phi(ab^{-1}))$ is a distance on G.
(2) For all a, b in G we have $d_\phi(a, b) = d_H(\phi(a), \phi(b))$.

Lemma 1 Condition (1) in the definition of a Gray map is equivalent to the following conditions,
(1) For $g \in G$ we have $w_H(\phi(g)) = 0$ iff $g = e$, where e stands for the identity of G,
(2) For all g in G we have $w_H(\phi(g)) = w_H(\phi(g^{-1}))$,
(3) For all x, y in G we have $w_H(\phi(xy)) \leq w_H(\phi(x)) + w_H(\phi(y))$.

Refer to in [1] for the proof of Lemma 1.

Example 2.1. Gray map over $C_4 = \langle x \mid x^4 = e \rangle$.

Example 2.2. Gray map over $K_4 = \langle x, y \mid x^2 = y^2 = e, xy = yx \rangle$. (We write K_4 for the *Klein four group.*) Gray maps as follows.

Example 2.1 and 2.2

C_4	$\mathbb{Z}_2^2$	K_4	$\mathbb{Z}_2^2$
$\phi(e)$	00	$\phi(e)$	00
$\phi(x)$	01	$\phi(x)$	01
$\phi(x^2)$	11	$\phi(y)$	11
$\phi(x^3)$	10	$\phi(xy)$	10

2.2 Type 1 Gray maps

In this subsection, we assume that H is a maximal subgroup of G with $[G : H] = 2$, and x is an arbitrary elements in $G \setminus H$ and h is an arbitrary elements in H. In this part, we construct a Gray map for G, called Type 1 Gray map, based on the existence of a Gray map for H. For $1 \in \mathbb{Z}_2$, let us denote by $\mathbf{1}$ the vector in $\mathbb{Z}_2^n$ whose components are all 1. Also we denote the usual concatenation of vectors in $\mathbb{Z}_2^n$ by $(\ |\ )$. Suppose $\phi : H \to \mathbb{Z}_2^n$ is a Gray map and define the map $\hat{\phi} : G \to \mathbb{Z}_2^{2n}$ by $\hat{\phi}(h) = (\phi(h) \mid \phi(h))$ and $\hat{\phi}(xh) = (\phi(h) \mid \phi(h) + \mathbf{1})$ in [1].

Lemma 2 For all $g \in G$ we have $w_H(\hat{\phi}(g)) = w_H(\hat{\phi}(g^{-1}))$.

Lemma 3 For all $a, b \in G$ we have $w_H(\hat{\phi}(ab)) \leq w_H(\hat{\phi}(a)) + w_H(\hat{\phi}(b))$.

Theorem 1 With notation as above, the map $\hat{\phi}$ is a Gray map.

Refer to in [1] for the proof of Lemma 2, 3 and Theorem 1.

2.3 Type 2 Gray maps

In this subsection, we assume that G is isomorphic to the semidirect product of two finite 2-groups H and K of orders 2^a and 2^b respectively, i.e. $G = H \ltimes_\psi K$ where $\psi : H \to \mathrm{Aut}(K)$ is a group homomorphism. Let $\theta_1 : H \to \mathbb{Z}_2^{n_1}$ and $\theta_2 : K \to \mathbb{Z}_2^{n_2}$ be Gray maps, where θ_2 is compatible with ψ in the sense that for all $h \in H$

$$w_H(\theta_2(k)) = w_H(\theta_2(\psi_h(k))).$$

Then we show that $\theta : G \to \mathbb{Z}_2^{n_1+n_2}$ with $\theta(hk) = (\theta_1(h) \mid \theta_2(k))$ is a Gray map, which we call Type 2 Gray map.

Theorem 2 With notation as above, if θ_2 is compatible with ψ, then the map θ is a Gray map.

Refer to in [1] for the proof of Theorem 2.

2.4 Automorphism

The set $\mathrm{Aut}(G)$ of all automorphisms of a group G forms a group under composition of mappings. Let X generate G. Each $\theta : G \to G$ in $\mathrm{Aut}(G)$ is determined by its values on X. For notational convenience we view the direct product of $C_4 = \langle x \rangle$ and $C_2 = \langle y \rangle$ as $\{\mathrm{e}, x, x^2, x^3, y, xy, x^2y, x^3y\}$. Because this group will appear frequently in this article we denote it by K_8 [2]. Similarly, $D_8 = \langle x, y \mid x^4 = y^2 = e, yx = x^3y \rangle$ is the dihedral group of order 8 and $Q_8 = \langle x, y \mid x^4 = e, y^2 = x^2, yx = x^3y \rangle$ is the quaternion group of order 8.

Example 2.3.

Aut(C_4) and Aut(C_8)$\simeq K_4$ [2]

$\mathrm{Aut}(C_4)$	effect on x	$\mathrm{Aut}(C_8)$	effect on x
φ_1	x	σ_1	x
φ_2	x^3	σ_2	x^3
		σ_3	x^5
		σ_4	x^7

Example 2.4.

Aut(K_8) $\simeq D_8$ [2]

$\mathrm{Aut}(K_8)$	effect on x	effect on y	order of automorphism
ψ_1	x	y	1
ψ_2	x^3y	x^2y	4
ψ_3	x^3	y	2
ψ_4	xy	x^2y	4
ψ_5	xy	y	2
ψ_6	x^3	x^2y	2
ψ_7	x^3y	y	2
ψ_8	x	x^2y	2

Example 2.5.

Aut(D_8)$\simeq D_8$

$\mathrm{Aut}(D_8)$	effect on x	effect on y	order of automorphism
α_1	x	y	1
α_2	x	xy	4
α_3	x	x^2y	2
α_4	x	x^3y	4
α_5	x^3	y	2
α_6	x^3	xy	2
α_7	x^3	x^2y	2
α_8	x^3	x^3y	2

3 Type 1 Gray maps for a group of order 8

Let G be $C_4 = \langle x \rangle$. Assume that $H = \{e, x^2\} \leq G$ be the maximal subgroup of G, which is isomorphic to C_2. Let $\phi_0 : H \to \mathbb{Z}_2$ be the natural map which sends e to 0 and x^2 to 1. Clearly ϕ_0 is a Gray map. Set $\phi_1 \stackrel{\text{def.}}{=} \hat{\phi}_0$, we have $\phi_1(e) = (\phi_0(e) \mid \phi_0(\mathrm{e})) = 00$, $\phi_1(x^2) = (\phi_0(x^2) \mid \phi_0(x^2)) = 11$, $\phi_1(x) = (\phi_0(e) \mid (\phi_0(\mathrm{e}) + 1)) = 01$ and $\phi_1(x^3) = (\phi_0(x^2) \mid (\phi_0(x^2) + 1)) = 10$. This is the well-known Gray map on C_4 in [1].

Example 3.1. Let G be $C_8 = \langle x \rangle$. Assume that $H = \{e, x^2, x^4, x^6\} \leq G$ be the maximal subgroup of G, which is isomorphic to C_4. Let $\phi_1 : H \to \mathbb{Z}_2^2$ be the previously constructed Gray map for C_4. We have,

$$\begin{array}{lll}
\hat{\phi}_1(e) & = (\phi_1(e) \mid \phi_1(e)) & = 0000 \\
\hat{\phi}_1(x^2) & = (\phi_1(x) \mid \phi_1(x)) & = 0101 \\
\hat{\phi}_1(x^4) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 1111 \\
\hat{\phi}_1(x^6) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 1010 \\
\hat{\phi}_1(x) & = (\phi_1(e) \mid \phi_1(e) + 11) & = 0011 \\
\hat{\phi}_1(x^3) & = (\phi_1(x) \mid \phi_1(x) + 11) & = 0110 \\
\hat{\phi}_1(x^5) & = (\phi_1(x^2) \mid \phi_1(x^2) + 11) & = 1100 \\
\hat{\phi}_1(x^7) & = (\phi_1(x^3) \mid \phi_1(x^3) + 11) & = 1001
\end{array}$$

Example 3.2. Let G be K_8. Similarly, assume that $H = \{e, x, x^2, x^3\} \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^2$ be the previously constructed Gray map for C_4. We have,

$$\begin{array}{lll}
\hat{\phi}_1(e) & = (\phi_1(e) \mid \phi_1(e)) & = 0000 \\
\hat{\phi}_1(x) & = (\phi_1(x) \mid \phi_1(x)) & = 0101 \\
\hat{\phi}_1(x^2) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 1111 \\
\hat{\phi}_1(x^3) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 1010 \\
\hat{\phi}_1(y) & = (\phi_1(e) \mid \phi_1(e) + 11) & = 0011 \\
\hat{\phi}_1(xy) & = (\phi_1(x) \mid \phi_1(x) + 11) & = 0110 \\
\hat{\phi}_1(x^2y) & = (\phi_1(x^2) \mid \phi_1(x^2) + 11) & = 1100 \\
\hat{\phi}_1(x^3y) & = (\phi_1(x^3) \mid \phi_1(x^3) + 11) & = 1001
\end{array}$$

Example 3.3. Let G be D_8. Similarly, assume that $H = \{e, x, x^2, x^3\} \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^2$ be the previously constructed Gray map for C_4. We have,

$$\begin{array}{lll}
\hat{\phi}_1(e) & = (\phi_1(e) \mid \phi_1(e)) & = 0000 \\
\hat{\phi}_1(x) & = (\phi_1(x) \mid \phi_1(x)) & = 0101 \\
\hat{\phi}_1(x^2) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 1111 \\
\hat{\phi}_1(x^3) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 1010 \\
\hat{\phi}_1(y) & = (\phi_1(e) \mid \phi_1(e) + 11) & = 0011 \\
\hat{\phi}_1(yx) & = (\phi_1(x) \mid \phi_1(x) + 11) & = 0110 \\
\hat{\phi}_1(yx^2) & = (\phi_1(x^2) \mid \phi_1(x^2) + 11) & = 1100 \\
\hat{\phi}_1(yx^3) & = (\phi_1(x^3) \mid \phi_1(x^3) + 11) & = 1001
\end{array}$$

Example 3.4. Let G be Q_8. Similarly, assume that $H = \{e, x, x^2, x^3\} \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^2$ be the previously constructed Gray map for C_4. We have,

$$\begin{array}{lll}
\hat{\phi}_1(e) & = (\phi_1(e) \mid \phi_1(e)) & = 0000 \\
\hat{\phi}_1(x) & = (\phi_1(x) \mid \phi_1(x)) & = 0101 \\
\hat{\phi}_1(x^2) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 1111 \\
\hat{\phi}_1(x^3) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 1010 \\
\hat{\phi}_1(y) & = (\phi_1(e) \mid \phi_1(e) + 11) & = 0011 \\
\hat{\phi}_1(yx) & = (\phi_1(x) \mid \phi_1(x) + 11) & = 0110 \\
\hat{\phi}_1(yx^2) & = (\phi_1(x^2) \mid \phi_1(x^2) + 11) & = 1100 \\
\hat{\phi}_1(yx^3) & = (\phi_1(x^3) \mid \phi_1(x^3) + 11) & = 1001
\end{array}$$

4 Type 2 Gray maps for a group of order 8

Example 4.1. Let G be D_8. We have $G \simeq C_2 \ltimes_\psi C_4$, where $\psi : C_2 \to \mathrm{Aut}(C_4)$ is a homomorphism. Consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_1 : C_4 \to \mathbb{Z}_2^2$ described in the example 2.1. Now we can construct a Type 2 Gray map θ from D_8 to $\mathbb{Z}_2^3$ as follows in [1],

$$\begin{array}{lll}
\theta(e) & = (\phi_0(e) \mid \phi_1(e)) & = 000 \\
\theta(x) & = (\phi_0(e) \mid \phi_1(x)) & = 001 \\
\theta(x^2) & = (\phi_0(e) \mid \phi_1(x^2)) & = 011 \\
\theta(x^3) & = (\phi_0(e) \mid \phi_1(x^3)) & = 010 \\
\theta(y) & = (\phi_0(y) \mid \phi_1(e)) & = 100 \\
\theta(yx) & = (\phi_0(y) \mid \phi_1(x)) & = 101 \\
\theta(yx^2) & = (\phi_0(y) \mid \phi_1(x^2)) & = 111 \\
\theta(yx^3) & = (\phi_0(y) \mid \phi_1(x^3)) & = 110
\end{array}$$

We denote by ϕ_2 the above Gray map θ.

Example 4.2. Similarly we can construct a Type 2 Gray map θ from K_8 to $\mathbb{Z}_2^3$,

$$\begin{array}{llll}
\theta(e) & = 000 & \theta(y) & = 100 \\
\theta(x) & = 001 & \theta(xy) & = 101 \\
\theta(x^2) & = 011 & \theta(x^2y) & = 111 \\
\theta(x^3) & = 010 & \theta(x^3y) & = 110
\end{array}$$

Also, we denote by ϕ_2 the above Gray map θ.

Since we have neither $C_8 \simeq C_2 \ltimes_\psi C_4$ nor $C_8 \simeq C_2 \ltimes_\psi K_4$, we can not construct a Type 2 Gray map for C_8.

Example 4.3. Let G be Q_8.

$$\begin{array}{llll}
\theta(e) & = 000 & \theta(y) & = 100 \\
\theta(x) & = 001 & \theta(yx) & = 101 \\
\theta(x^2) & = 011 & \theta(yx^2) & = 111 \\
\theta(x^3) & = 010 & \theta(yx^3) & = 110
\end{array}$$

In Q_8, we have $x^{-1} = x^3, y^{-1} = yx^2, (yx)^{-1} = yx^3$, so we have $w_H(\phi(g)) \neq w_H(\phi(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for Q_8.

In $\mathbb{Z}_2^3$ weight distribution is as follows:

Hamming-weight	0	1	2	3
number of points	1	3	3	1

Since both 1 and 3 are odd, in order to satisfy (2) of Lemma 1, the group G must have at least 3 elements of order 2.

On the other hand, C_8 has one element of order 2, therefore can not satisfy (2) of Lemma 1. Similarly, neither does Q_8.

5 Type 1 Gray map for a group of order 16

5.1 Cyclic extensions.

We state the classification of groups of order 16. Let $N \lhd G$ and $N' \lhd G'$. Suppose that $G/N \simeq C_n$.(Such a group G is called an *(inner) cyclic extension* of N.) Pick any a in $G \setminus N$ such that the coset Na has order n in G/N. Then $v = a^n$ is in N, and n is minimal with that property. Further, let τ in $\mathrm{Aut}(N)$ be the restriction of $t_a(x) \stackrel{\text{def.}}{=} axa^{-1}$ to N. Then

$\tau(v) = aa^n a^{-1} = a^n = v$, and
$\tau^n = a \cdots a(axa^{-1})a^{-1} \cdots a^{-1} = a^n x a^n = vxv^{-1} = t_v(x)$, for all x in N,
so $\tau^n = t_v$.

Definition 4 A quadruple (N, n, τ, v) is an *extension type* if N is a group and if v in N and τ in $\mathrm{Aut}(N)$ are such that $\tau(v) = v$ and $\tau^n = t_v$.

Definition 5 The extension types (N, n, τ, v) and (N', n, σ, w) are *equivalent* if there is an isomorphism $\phi : N \to N'$ such that $\sigma = \phi \circ \tau \circ \phi^{-1}$ and $w = \phi(v)$ in [2].

Theorem 3 There are exactly 14 groups of order 16 up to isomorphism. Besides the outsider $G_0 = C_2 \times C_2 \times C_2 \times C_2$ they can be listed as follows,

$$
\begin{array}{ll}
G_1 = C_2 \times C_8 & (C_8, 2, \sigma_1, e) \\
G_2 = C_2 \ltimes_{\sigma_2} C_8 & (C_8, 2, \sigma_2, e) \\
G_3 = C_2 \ltimes_{\sigma_3} C_8 & (C_8, 2, \sigma_3, e) \\
G_4 = C_2 \ltimes_{\sigma_4} C_8 & (C_8, 2, \sigma_4, e) \\
G_5 = Q_{16} & (C_8, 2, \sigma_4, x^4) \\
G_6 = C_{16} & (C_8, 2, \sigma_1, x) \\
G_7 = K_4 \times C_4 & (K_8, 2, \psi_1, e) \\
G_8 = D_8 \times C_2 & (K_8, 2, \psi_3, e) \\
G_9 = C_4 \ltimes_{\tau} K_4 & (K_8, 2, \psi_5, e) \\
G_{10} = C_2 \ltimes_{\tau} Q_8 & (K_8, 2, \psi_6, e) \\
G_{11} = C_2 \times Q_8 & (K_8, 2, \psi_3, x^2) \\
G_{12} = C_4 \ltimes_{\tau} C_4 & (K_8, 2, \psi_5, x^2) \\
G_{13} = C_4 \times C_4 & (K_8, 2, \psi_1, y)
\end{array}
$$

Refer to in [2] for the proof of Theorem 3.

5.2 Type 1 Gray map for G_1.

Example 5.1. Let G be $G_1(C_8, 2, \sigma_1, e)$. Assume that $H = C_8 \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^4$ be the previously constructed Gray map for C_8. Set $\phi_2 \stackrel{\text{def.}}{=} \hat{\phi}_1$, we have,

$$
\begin{array}{lll}
\phi_2(e) & = (\phi_1(e) \mid \phi_1(e)) & = 00000000 \\
\phi_2(x) & = (\phi_1(x) \mid \phi_1(x)) & = 00110011 \\
\phi_2(x^2) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 01010101 \\
\phi_2(x^3) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 01100110 \\
\phi_2(x^4) & = (\phi_1(x^4) \mid \phi_1(x^4)) & = 11111111 \\
\phi_2(x^5) & = (\phi_1(x^5) \mid \phi_1(x^5)) & = 11001100 \\
\phi_2(x^6) & = (\phi_1(x^6) \mid \phi_1(x^6)) & = 10101010 \\
\phi_2(x^7) & = (\phi_1(x^7) \mid \phi_1(x^7)) & = 10011001 \\
\phi_2(a) & = (\phi_1(e) \mid \phi_1(e) + 1111) & = 00001111 \\
\phi_2(xa) & = (\phi_1(x) \mid \phi_1(x) + 1111) & = 00111100 \\
\phi_2(x^2a) & = (\phi_1(x^2) \mid \phi_1(x^2) + 1111) & = 01011010 \\
\phi_2(x^3a) & = (\phi_1(x^3) \mid \phi_1(x^3) + 1111) & = 01101001 \\
\phi_2(x^4a) & = (\phi_1(x^4) \mid \phi_1(x^4) + 1111) & = 11110000 \\
\phi_2(x^5a) & = (\phi_1(x^5) \mid \phi_1(x^5) + 1111) & = 11000011 \\
\phi_2(x^6a) & = (\phi_1(x^6) \mid \phi_1(x^6) + 1111) & = 10100101 \\
\phi_2(x^7a) & = (\phi_1(x^7) \mid \phi_1(x^7) + 1111) & = 10010110
\end{array}
$$

5.3 Type 1 Gray map for G_2, G_3, G_4, G_5 and G_6.

Let G be $G_2(C_8, 2, \sigma_2, e)$. Assume that $H = C_8 \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^4$ be the previously constructed Gray map for C_8. Set $\phi_2 \stackrel{\text{def.}}{=} \hat{\phi}_1$, we have the same Gray map with Example 5.1 for G_2. Also we have the same Gray map with Example 5.1 for G_3, G_4, G_5 and G_6.

5.4 Type 1 Gray map for G_7.

Example 5.2. Let G be $G_7(K_8, 2, \psi_1, e)$. Assume that $H = K_8 \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^4$ be the previously constructed Gray map for K_8. Set $\phi_2 \stackrel{\text{def.}}{=} \hat{\phi}_1$, we have,

$$\begin{array}{lll}
\phi_2(e) & = (\phi_1(e) \mid \phi_1(e)) & = 00000000 \\
\phi_2(x) & = (\phi_1(x) \mid \phi_1(x)) & = 00110011 \\
\phi_2(x^2) & = (\phi_1(x^2) \mid \phi_1(x^2)) & = 01010101 \\
\phi_2(x^3) & = (\phi_1(x^3) \mid \phi_1(x^3)) & = 01100110 \\
\phi_2(y) & = (\phi_1(y) \mid \phi_1(y)) & = 11111111 \\
\phi_2(xy) & = (\phi_1(xy) \mid \phi_1(xy)) & = 11001100 \\
\phi_2(x^2y) & = (\phi_1(x^2y) \mid \phi_1(x^2y)) & = 10101010 \\
\phi_2(x^3y) & = (\phi_1(x^3y) \mid \phi_1(x^3y)) & = 10011001 \\
\phi_2(a) & = (\phi_1(e) \mid (\phi_1(e) + 1111)) & = 00001111 \\
\phi_2(xa) & = (\phi_1(x) \mid (\phi_1(x) + 1111)) & = 00111100 \\
\phi_2(x^2a) & = (\phi_1(x^2) \mid (\phi_1(x^2) + 1111)) & = 01011010 \\
\phi_2(x^3a) & = (\phi_1(x^3) \mid (\phi_1(x^3) + 1111)) & = 01101001 \\
\phi_2(ya) & = (\phi_1(y) \mid (\phi_1(y) + 1111)) & = 11110000 \\
\phi_2(xya) & = (\phi_1(xy) \mid (\phi_1(xy) + 1111)) & = 11000011 \\
\phi_2(x^2ya) & = (\phi_1(x^2y) \mid (\phi_1(x^2y) + 1111)) & = 10100101 \\
\phi_2(x^3ya) & = (\phi_1(x^3y) \mid (\phi_1(x^3y) + 1111)) & = 10010110
\end{array}$$

5.5 Type 1 Gray map for $G_8, G_9, G_{10}, G_{11}, G_{12}$ and G_{13}.

Let G be $G_8(K_8, 2, \psi_3, e)$. Assume that $H = K_8 \leq G$ be the maximal subgroup of G. Let $\phi_1 : H \to \mathbb{Z}_2^4$ be the previously constructed Gray map for K_8. Set $\phi_2 \stackrel{\text{def.}}{=} \hat{\phi}_1$, we have the same Gray map with Example 5.2 for G_8. Also we have the same Gray map with Example 5.2 for $G_9, G_{10}, G_{11}, G_{12}$ and G_{13}.

6 Type 2 Gray maps for a group of order 16

6.1 Type 2 Gray map for G_1,G_2, G_3, G_4, G_5 and G_6.

We can not construct a Type 2 Gray map for G_1,G_2, G_3, G_4, G_5 and G_6 which are semidirect products of C_2 and C_8, because we can not construct a Type 2 Gray map for C_8. (Section 4)

6.2 Type 2 Gray map for G_7.

Example 6.1. Let G be G_7. We have $G \simeq C_2 \times K_8$ and $v = a^2 = e$. Consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the example 4.2. Now we can construct a Type 2 Gray map θ from G_7 to $\mathbb{Z}_2^4$ as follows,

$$
\begin{array}{lll}
\theta(e) & = (\phi_0(e) \mid \phi_2(e)) & = 0000 \\
\theta(x) & = (\phi_0(e) \mid \phi_2(x)) & = 0001 \\
\theta(x^2) & = (\phi_0(e) \mid \phi_2(x^2)) & = 0011 \\
\theta(x^3) & = (\phi_0(e) \mid \phi_2(x^3)) & = 0010 \\
\theta(y) & = (\phi_0(e) \mid \phi_2(y)) & = 0100 \\
\theta(xy) & = (\phi_0(e) \mid \phi_2(xy)) & = 0101 \\
\theta(x^2y) & = (\phi_0(e) \mid \phi_2(x^2y)) & = 0111 \\
\theta(x^3y) & = (\phi_0(e) \mid \phi_2(x^3y)) & = 0110 \\
\theta(a) & = (\phi_0(a) \mid \phi_2(e)) & = 1000 \\
\theta(xa) & = (\phi_0(a) \mid \phi_2(x)) & = 1001 \\
\theta(x^2a) & = (\phi_0(a) \mid \phi_2(x^2)) & = 1011 \\
\theta(x^3a) & = (\phi_0(a) \mid \phi_2(x^3)) & = 1010 \\
\theta(ya) & = (\phi_0(a) \mid \phi_2(y)) & = 1100 \\
\theta(xya) & = (\phi_0(a) \mid \phi_2(xy)) & = 1101 \\
\theta(x^2ya) & = (\phi_0(a) \mid \phi_2(x^2y)) & = 1111 \\
\theta(x^3ya) & = (\phi_0(a) \mid \phi_2(x^3y)) & = 1110
\end{array}
$$

In G_7, we have $(xa)^{-1} = x^3a, (xya)^{-1} = x^3ya$. So (2) and (3) of Lemma 1 hold. Since in $G_7 \simeq C_2 \times K_8$, we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all h $\in C_2(= \{e, a\})$, and therefore θ is a Type 2 Gray map for G_7 by Theorem 2. We can construct a Type 2 Gray map for G_7.

6.3 Type 2 Gray map for G_8.

Let G be G_8. We have $G \simeq C_2 \ltimes_{\psi_3} K_8$ and $v = a^2 = e$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct a similar Type 2 Gray map as in Example 6.1. The orders of all elements in $G_8 \setminus K_8$ are 2, so (2) and (3) of Lemma 1 hold. Since in $G_8 \simeq C_2 \ltimes_{\psi_3} K_8$, where $ax = x^3a$ and $ay = ya$, we have $w_H(\phi_2(\psi_a(x))) = w_H(\phi_2(axa^{-1}) = w_H(\phi_2(x^3) = w_H(\phi_2(x))$ and $w_H(\phi_2(\psi_a(xy))) = w_H(\phi_2(axya^{-1}) = w_H(\phi_2(x^3y) = w_H(\phi_2(xy))$ and so on. Hence we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all $h \in C_2(= \{e, a\})$, and therefore θ is a Type 2 Gray map for G_8 by Theorem 2. We can construct a Type 2 Gray map for G_8.

6.4 Type 2 Gray map for $G_8(D_8, 2, \alpha_5, e)$

In [6], D_8 is one of the maximal subgroups of G_8. So, we consider to construct a Type 2 Gray map for G_8 with $N = D_8$ instead of $N = K_8$.

Let G be $G_8(D_8, 2, \alpha_5, e)$. We have $G \simeq C_2 \ltimes_{\alpha_5} D_8$. Consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : D_8 \to \mathbb{Z}_2^3$ described in the Example 4.1. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_8 we have $(xya)^{-1} = x^3ya$, so (2) and (3) of Lemma 1 hold. Since in $G_8 \simeq C_2 \ltimes_{\alpha_5} D_8$, where $ax = x^3a$ and $ay = ya$, so we have $w_H(\phi_2(\psi_a(x))) = w_H(\phi_2(axa^{-1}) = w_H(\phi_2(x^3) = w_H(\phi_2(x))$ and $w_H(\phi_2(\psi_a(xy))) = w_H(\phi_2(axya^{-1}) = w_H(\phi_2(x^3y) = w_H(\phi_2(xy))$ and so on. Hence we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all h $\in C_2(= \{e, a\})$, and therefore θ is a Type 2 Gray map for G_8 by Theorem 2. We can construct a Type 2 Gray map for G_8.

6.5 Type 2 Gray map for G_9.

Let G be G_9. We have $G \simeq C_2 \ltimes_{\psi_5} K_8$ and $v = a^2 = e$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_9 we have $(xa)^{-1} = x^3ya$ and $(x^3a)^{-1} = xya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_9.

6.6 Type 2 Gray map for G_{10}.

Let G be G_{10}. We have $G \simeq C_2 \ltimes_{\psi_6} K_8$ and $v = a^2 = e$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct the a similar Type 2 Gray map as in Example 6.1. In G_{10} we have $(xy)^{-1} = x^3y, (ya)^{-1} = x^2ya$ and $(xya)^{-1} = x^3ya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_{10}.

6.7 Type 2 Gray map for $G_{10}(D_8, 2, \alpha_3, e)$

In [6], D_8 is one of the maximal subgroups of G_{10}. So we consider to construct a Type 2 Gray map for G_{10} with $N = D_8$ instead of $N = K_8$.

Let G be $G_{10}(D_8, 2, \alpha_3, e)$. We have $G \simeq C_2 \ltimes_{\alpha_3} D_8$. Consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : D_8 \to \mathbb{Z}_2^3$ described in the Example 4.1. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_8 we have $(xa)^{-1} = x^3a, (ya)^{-1} = x^2ya$ and $(xya)^{-1} = x^3ya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_{10}.

6.8 Type 2 Gray map for G_{11}.

Let G be G_{11}. We have $G \simeq C_2 \ltimes_{\psi_3} K_8$ and $v = a^2 = x^2$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_{11} we have $a^{-1} = x^2a, (xa)^{-1} = x^3a, (ya)^{-1} = x^2ya$ and $(xya)^{-1} = x^3ya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_{11}.

6.9 Type 2 Gray map for G_{12}.

Let G be G_{12}. We have $G \simeq C_2 \ltimes_{\psi_5} K_8$ and $v = a^2 = x^2$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_{12} we have $a^{-1} = x^2a, (xa)^{-1} = xya, (x^3a)^{-1} = x^3ya$ and $(ya)^{-1} = x^2ya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_{12}.

6.10 Type 2 Gray map for G_{13}.

Let G be G_{13}. We have $G \simeq C_2 \ltimes_{\psi_1} K_8$ and $v = a^2 = y$. Similarly, consider the natural Gray map $\phi_0 : C_2 \to \mathbb{Z}_2$ and the Gray map $\phi_2 : K_8 \to \mathbb{Z}_2^3$ described in the Example 4.2. Now we can construct a similar Type 2 Gray map as in Example 6.1. In G_{13} we have $a^{-1} = ya, (xa)^{-1} = x^3ya, (x^2a)^{-1} = x^2ya$ and $(x^3a)^{-1} = xya$, so we have $w_H(\theta(g)) \neq w_H(\theta(g^{-1}))$. Since θ does not satisfy (2) of Lemma 1, we can not construct a Type 2 Gray map for G_{13}.

7 The extension type $(C_4, 4, \tau, v)$ or $(K_4, 4, \tau, v)$

In Theorem 3 we have $G_7 = C_4 \times K_4$, $G_9 = C_4 \ltimes_\tau K_4$, $G_{12} = C_4 \ltimes_\tau C_4$ and $G_{13} = C_4 \times C_4$, so we consider to construct a Type 2 Gray map for G with $N = C_4$ or $N = K_4$.

7.1 Type 2 Gray map for $G_7(K_4, 4, \phi_1, e)$

We assume that K_4 is $\langle a\rangle \times \langle y\rangle$ and C_4 is $\langle x\rangle$, where $x^4 = y^2 = a^2 = e$. Let G be G_7. We have $G \simeq C_4 \times K_4$ and $v = x^4 = e$. Consider the Gray map $\phi_1 : C_4 \to \mathbb{Z}_2^2$ and the Gray map $\phi_2 : K_4 \to \mathbb{Z}_2^2$ described in the example 2.1 and 2.2. Now we can construct a Type 2 Gray map θ from G_7 to $\mathbb{Z}_2^4$ as follows,

$$\begin{array}{lll}
\theta(e) & = (\phi_1(e) \mid \phi_2(e)) & = 0000 \\
\theta(a) & = (\phi_1(e) \mid \phi_2(a)) & = 0001 \\
\theta(y) & = (\phi_1(e) \mid \phi_2(y)) & = 0011 \\
\theta(ya) & = (\phi_1(e) \mid \phi_2(ya)) & = 0010 \\
\theta(x) & = (\phi_1(x) \mid \phi_2(e)) & = 0100 \\
\theta(xa) & = (\phi_1(x) \mid \phi_2(a)) & = 0101 \\
\theta(xy) & = (\phi_1(x) \mid \phi_2(y)) & = 0111 \\
\theta(xya) & = (\phi_1(x) \mid \phi_2(ya)) & = 0110 \\
\theta(x^2) & = (\phi_1(x^2) \mid \phi_2(e)) & = 1100 \\
\theta(x^2a) & = (\phi_1(x^2) \mid \phi_2(a)) & = 1101 \\
\theta(x^2y) & = (\phi_1(x^2) \mid \phi_2(y)) & = 1111 \\
\theta(x^2ya) & = (\phi_1(x^2) \mid \phi_2(ya)) & = 1110 \\
\theta(x^3) & = (\phi_1(x^3) \mid \phi_2(e)) & = 1000 \\
\theta(x^3a) & = (\phi_1(x^3) \mid \phi_2(a)) & = 1001 \\
\theta(x^3y) & = (\phi_1(x^3) \mid \phi_2(y)) & = 1011 \\
\theta(x^3ya) & = (\phi_1(x^3) \mid \phi_2(ya)) & = 1010
\end{array}$$

In G_7 we have $(xa)^{-1} = x^3a, (xy)^{-1} = x^3y$ and $(xya)^{-1} = x^3ya$, so (2) and (3) of Lemma 1 hold. Since in $G_7 \simeq C_4 \times K_4$, we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all h $\in C_4 (= \{e, x, x^2, x^3\})$, and therefore θ is a Type 2 Gray map for G_7 by Theorem 2. We can construct a Type 2 Gray map for G_7.

7.2 Type 2 Gray map for $G_9(K_4, 4, \sigma, e)$

In [2], $G_9 \simeq \langle x\rangle \ltimes_\tau \langle y, a\rangle = C_4 \ltimes_\tau K_4$, where $\tau(y) = y$ and $\tau(a) = ya$ (i.e. $x(y)x^{-1} = y$ and $x(a)x^{-1} = ya$). Because all order two elements σ of $\mathrm{Aut}(K_4) \simeq S_3$ are conjugate, they yield isomorphic semidirect products $C_4 \ltimes_\sigma K_4$.

Let G be G_9. We have $G \simeq C_4 \ltimes_\sigma K_4$ and $v = x^4 = e$. Consider the Gray map $\phi_1 : C_4 \to \mathbb{Z}_2^2$ and the Gray map $\phi_2 : K_4 \to \mathbb{Z}_2^2$ described in the Example 2.1 and 2.2. Now we can construct a Type 2 Gray map θ from G_9 to $\mathbb{Z}_2^4$ as follows,

$$
\begin{array}{lll}
\theta(e) & = (\phi_1(e) \mid \phi_2(e)) & = 0000 \\
\theta(a) & = (\phi_1(e) \mid \phi_2(a)) & = 0001 \\
\theta(y) & = (\phi_1(e) \mid \phi_2(y)) & = 0011 \\
\theta(ya) & = (\phi_1(e) \mid \phi_2(ya)) & = 0010 \\
\theta(x) & = (\phi_1(x) \mid \phi_2(e)) & = 0100 \\
\theta(xa) & = (\phi_1(x) \mid \phi_2(a)) & = 0101 \\
\theta(xy) & = (\phi_1(x) \mid \phi_2(y)) & = 0111 \\
\theta(xya) & = (\phi_1(x) \mid \phi_2(ya)) & = 0110 \\
\theta(x^2) & = (\phi_1(x^2) \mid \phi_2(e)) & = 1100 \\
\theta(x^2a) & = (\phi_1(x^2) \mid \phi_2(a)) & = 1101 \\
\theta(x^2y) & = (\phi_1(x^2) \mid \phi_2(y)) & = 1111 \\
\theta(x^2ya) & = (\phi_1(x^2) \mid \phi_2(ya)) & = 1110 \\
\theta(x^3) & = (\phi_1(x^3) \mid \phi_2(e)) & = 1000 \\
\theta(x^3a) & = (\phi_1(x^3) \mid \phi_2(a)) & = 1001 \\
\theta(x^3y) & = (\phi_1(x^3) \mid \phi_2(y)) & = 1011 \\
\theta(x^3ya) & = (\phi_1(x^3) \mid \phi_2(ya)) & = 1010
\end{array}
$$

In G_9 we have $(xy)^{-1} = x^3y, (xa)^{-1} = x^3ya$ and $(x^3a)^{-1} = xya$, so (2) and (3) of Lemma 1 hold. Since $G_9 \simeq C_4 \ltimes_\sigma K_4$, where $xy = yx$ and $xa = yax$, so we have $w_H(\phi_2(\psi_x(a))) = w_H(\phi_2(xax^{-1}) = w_H(\phi_2(ya) = w_H(\phi_2(a))$ and $w_H(\phi_2(\psi_{x^2}(a))) = w_H(\phi_2(x^2ax^2) = w_H(\phi_2(a))$ and so on. Hence we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all h $\in C_4(= \{e, x, x^2, x^3\})$, and therefore θ is a Type 2 Gray map for G_9 by Theorem 2. We can construct a Type 2 Gray map for G_9.

7.3 Type 2 Gray map for $G_{12}(C_4, 4, \varphi_2, e)$

Similarly $G_{12} \simeq \langle x \rangle \ltimes_\tau \langle xa \rangle = C_4 \ltimes_\tau C_4$, where $\langle xa \rangle = \{e, xa, y, xya\}, \tau = \varphi_2$ (i.e. $\varphi_2(xa) = xya$) and $v = a^4 = e$. Consider the Gray map $\phi_1 : C_4(= \langle x \rangle) \to \mathbb{Z}_2^2$ and the Gray map $\phi_2 : C_4(= \langle xa \rangle) \to \mathbb{Z}_2^2$ described in the Example 2.1. Now we can construct a Type 2 Gray map θ from G_{12} to $\mathbb{Z}_2^4$ as follows,

$$
\begin{array}{lll}
\theta(e) & = (\phi_1(e) \mid \phi_2(e)) & = 0000 \\
\theta(xa) & = (\phi_1(e) \mid \phi_2(xa)) & = 0001 \\
\theta(y) & = (\phi_1(e) \mid \phi_2(y)) & = 0011 \\
\theta(xya) & = (\phi_1(e) \mid \phi_2(xya)) & = 0010 \\
\theta(x) & = (\phi_1(x) \mid \phi_2(e)) & = 0100 \\
\theta(x^2a) & = (\phi_1(x) \mid \phi_2(xa)) & = 0101 \\
\theta(xy) & = (\phi_1(x) \mid \phi_2(y)) & = 0111 \\
\theta(x^2ya) & = (\phi_1(x) \mid \phi_2(xya)) & = 0110 \\
\theta(x^2) & = (\phi_1(x^2) \mid \phi_2(e)) & = 1100 \\
\theta(x^3a) & = (\phi_1(x^2) \mid \phi_2(xa)) & = 1101 \\
\theta(x^2y) & = (\phi_1(x^2) \mid \phi_2(y)) & = 1111 \\
\theta(x^3ya) & = (\phi_1(x^2) \mid \phi_2(xya)) & = 1110 \\
\theta(x^3) & = (\phi_1(x^3) \mid \phi_2(e)) & = 1000 \\
\theta(a) & = (\phi_1(x^3) \mid \phi_2(xa)) & = 1001 \\
\theta(x^3y) & = (\phi_1(x^3) \mid \phi_2(y)) & = 1011 \\
\theta(ya) & = (\phi_1(x^3) \mid \phi_2(xya)) & = 1010
\end{array}
$$

In G_{12} we have $(xy)^{-1} = x^3y, a^{-1} = x^2a, (xa)^{-1} = xya, x^3a^{-1} = x^3ya$ and $(ya)^{-1} = x^2ya$, so (2) and (3) of Lemma 1 hold. Since in $G_{12} \simeq C_4 \ltimes_{\varphi_2} C_4$, where $\varphi_2(xa) = xya$, so we have $w_H(\phi_2(\psi_x(xa))) = w_H(\phi_2(xya) = w_H(\phi_2(xa)$ and so on. Hence we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all $h \in C_4(= \{e, x, x^2, x^3\})$, and therefore θ is a Type 2 Gray map for G_{12} by Theorem 2. We can construct a Type 2 Gray map for G_{12}.

7.4 Type 2 Gray map for $G_{13}(C_4, 4, \varphi_1, e)$

In [2], $G_{13} \simeq C_4(= \langle x \rangle) \times C_4(= \langle a \rangle)$, where $\langle a \rangle = \{e, a, y, ya\}, \tau(a) = \varphi_1(a) = a$ (namely $xa = ax$) and $v = a^4 = y^2 = e$. Consider the Gray map $\phi_1 : C_4(= \langle x \rangle) \to \mathbb{Z}_2^2$ and the Gray map $\phi_2 : C_4(= \langle a \rangle) \to \mathbb{Z}_2^2$ described in the example 2.1. Now we can construct a Type 2 Gray map θ from G_{13} to $\mathbb{Z}_2^4$ as follows,

$$\begin{array}{lll}
\theta(e) & = (\phi_1(e) \mid \phi_2(e)) & = 0000 \\
\theta(a) & = (\phi_1(e) \mid \phi_2(a)) & = 0001 \\
\theta(y) & = (\phi_1(e) \mid \phi_2(y)) & = 0011 \\
\theta(ya) & = (\phi_1(e) \mid \phi_2(ya)) & = 0010 \\
\theta(x) & = (\phi_1(x) \mid \phi_2(e)) & = 0100 \\
\theta(xa) & = (\phi_1(x) \mid \phi_2(a)) & = 0101 \\
\theta(xy) & = (\phi_1(x) \mid \phi_2(y)) & = 0111 \\
\theta(xya) & = (\phi_1(x) \mid \phi_2(ya)) & = 0110 \\
\theta(x^2) & = (\phi_1(x^2) \mid \phi_2(e)) & = 1100 \\
\theta(x^2a) & = (\phi_1(x^2) \mid \phi_2(a)) & = 1101 \\
\theta(x^2y) & = (\phi_1(x^2) \mid \phi_2(y)) & = 1111 \\
\theta(x^2ya) & = (\phi_1(x^2) \mid \phi_2(ya)) & = 1110 \\
\theta(x^3) & = (\phi_1(x^3) \mid \phi_2(e)) & = 1000 \\
\theta(x^3a) & = (\phi_1(x^3) \mid \phi_2(a)) & = 1001 \\
\theta(x^3y) & = (\phi_1(x^3) \mid \phi_2(y)) & = 1011 \\
\theta(x^3ya) & = (\phi_1(x^3) \mid \phi_2(ya)) & = 1010
\end{array}$$

In G_{13}, we have $a^{-1} = ya, x^{-1} = x^3, (xa)^{-1} = x^3ya, (xy)^{-1} = x^3y, (xya)^{-1} = x^3a$ and $(x^2a)^{-1} = x^2ya$, so (2) and (3) of Lemma 1 hold. Since in $G_{13} \simeq C_4 \times C_4$, we have $w_H(\phi_2(k)) = w_H(\phi_2(\psi_h(k)))$ for all h $\in C_4(= \{e, x, x^2, x^3\})$, and therefore θ is a Type 2 Gray map for G_{13} by Theorem 2. we can construct a Type 2 Gray map for G_{13}.

8 Summary

8.1 Type 1 Gray map

We can construct a Type 1 Gray map for $G_0, G_1, \ldots, G_{12}$ and G_{13}.

8.2 Type 2 Gray map

1. We can construct a Type 2 Gray map for $G_0, G_7, G_8, G_9, G_{12}$ and G_{13}.
2. A necessary conditions that we can construct a Type 2 Gray map for a group of order 16 are,
(1) In the extension type $(N, 2, \tau, v)$ N is K_8 or D_8 or $C_2 \times C_2 \times C_2$ and v is e.
(2) Alternatively in the extension type $(N, 4, \tau, v)$ N is C_4 or K_4 and v is e.
(3) G do not contain the subgroup C_8 or Q_8.

References

1. Reza Sobhani, Gray isometries for finite p-groups, Transactions on Combinatorics, Vol.2 No.1(2013), pp17-26.

2. Marcel Wild, The groups of order sixteen made easy, The Mathematical Association of America, January 2005.

3. A. R. Hammons, P. V. Kummar, A. R. Calderbank, N. J. A. Sloane and P. Sole, The $\mathbb{Z}_4$ linearity of Kerdock, Preparata, Goethals and related codes, IEEE Trans. Inform. Theory, 40(1994).

4. Joseph J. Rotman, An introduction to the Theory of Groups, Springer 1995.

5. J. H. van Lint, Introduction to Coding Theory, Third Edition, Springer 1999.

6. A. D. Thomas and G. V. Wood, Group Tables, Mathematics Series 2, Shiva Publishing Limited, Kent, UK, 1980.

7. F. J. Macwilliams and N. J. A. Sloane, The theory of error-correcting codes, North-Holland, Amsterdam, 1977.